

Garlic Rust (*Puccinia Allii*): Effect and Management Options- A Review

Tilahun Negash¹, Hassen Shifa²

Summary

Garlic rust caused by *Puccinia allii* is potentially damaging to garlic crops, has a wide distribution and limits production of the crop in many countries. Different disease management options are being used to reduce its effect. Cultural disease management methods such as rotation with non-host crops, optimum crop density, adequate supply of moisture, use of healthy planting materials and field sanitation with other crops can reduce the intensity of garlic rust by reducing available inoculum to initiate infection. Use of host plant resistance to manage diseases reported as it is economical, long-lasting, effective, easy to handle and environment-friendly. There are several fungicides which are effective against garlic rust, as a stop-gap measure or as an integral part of the crop management system. Integrated management of garlic rust is possible that a combination of control tactics is applied in many parts of the world. Such an integrated approach does not depend on a single method, which in the case of monogenic host resistance could be non-durable, and should be more sustainable over time.

Keywords: *Allium sativum* L., *Puccinia allii*, Effect, Management

Introduction

Garlic (*Allium sativum* L.) Garlic plant is principally used as spice for flavoring and seasoning vegetable and meat dishes, it gives the food a delightful fragrance. In addition, garlic has miracle pharmaceutical effects and used to cure an enormous disease attributed to its rich content of sulfur-containing compounds, i.e., alliin, g-glutamylcysteine, and their derivatives (Chia-Wen Tsai *et al.*, 2012).

World garlic cultivation increased from 1,204,711 ha of land in 2007 with total production from 6.5 to 15.68 million tons to 1,199,929 ha with a production of 17674893 tonnes (FAO, 2012). In Ethiopia, 15,381.01 ha of land was under garlic cultivation with a production of about 6.15 tones/ ha (CSA, 2017). It is the second most widely cultivated *Allium* species next to onion in Ethiopia (Zewde *et al.*, 2007). Garlic is produced mainly in the mid and highlands of Ethiopia (Getachew and Asfaw, 2000). Garlic can be grown under a wide range of climatic conditions but prefers cool weather and grows at higher elevation (900 to 1200 meters) and grows best within the geographic areas having a mean monthly growing temperature ranging from 12oC to 24oC (Libner, 1989). In most areas elevation from 500-2000 m provide suitable growing conditions, particularly during dry periods (Tindall, 1983). It had also been under commercial production by Horticultural Development Corporation at Debre Zeit, Guder and Tseday State Farms (Getachew and Asfaw, 2000). The crop is produced mainly as a cash crop to earn foreign currency by exporting it to Europe, the Middle East and USA.

Garlic production has been significantly reduced due to fungal disease. Among the fungal diseases, foliar disease, caused by the fungus are some of the most destructive plant diseases worldwide (Woldeab *et al.*, 2007; Tome *et al.*, 2010). From foliar disease garlic rust caused by *Puccinia allii* is the major destructive disease problem in almost all garlic producing regions of Ethiopia (Tesfaye and Habtu, 2003; Mengesha *et al.*, 2016). *Puccinia allii* is an autoecious rust, producing pycnia, aecia, uredinia, telia, and basidia on a single host (Anikster *et al.*, 2004). The genus *Puccinia* is the largest in the order Uredinales. *Puccinia allii* infects garlic at bulb formation stage (Koike *et al.*, 2001).

¹ Department of Plant Science, College of Agriculture and Natural Resource Management, Bonga University, P.O. Box 334, Bonga, Ethiopia. E-mail: tilatanejash211@gmail.com

² Department of Plant Science, College of Agriculture and Natural Resource, Madda Walabu University, P.O. Box 247, Bale Robe, Ethiopia

Garlic rust is readily identified by the earliest symptom of small, circular to elongate white flecks that occur on both sides of leaves, as the disease progresses, these small spots expand, and the leaf tissue covering the lesions ruptures and masses of orange, powdery spores (uredospores) become visible as pustules (Schwartz and Mohan, 2008).

The disease has been found in every parts of Ethiopia where garlic are cultivated (Tesfaye and Habtu, 2003; Mengesha *et al.*, 2016; Worku, 2017). The disease does not attack the garlic bulb directly, but it damages on the leaves which indirect effect size and quality of bulbs at harvest thereby reduce its marketability (Koike *et al.*, 2001, Tahir *et al.*, 2006). Until recently, this fungus was considered to be of minor importance in garlic production. However, outbreaks in USA in the late 1990's reduced crop yields by up to 75% in some fields (Anikster *et al.*, 2004). The disease also occurred in 1999 and 2000, indicating that garlic rust may have become an annual problem in some parts of the United States of America (Koike *et al.*, 2001). Heavily infected plants may be more susceptible to secondary infections and there can be direct bulb yield losses.

Worldwide, garlic rust has caused significant losses to garlic, leek, and onion production. Yield losses as high as 83% have also been reported by Ahmad and Iqbal (2002) on garlic due to rust disease in Nepal. Bulbs infected with garlic rust remain small and are of low storage quality (Sartaj and Ahmad, 2005). Garlic rust is the most important disease of the crop in Ethiopia causing a total bulb yield loss as high as 58.75% (Worku, 2017). Because the disease is destructive, different management options are being used to reduce effect of rust to garlic. Cultural disease management methods such as rotation with non-host crops, field sanitation, late planting and intercropping of garlic with other crops can reduce the incidence and severity of garlic rust, by reducing available inoculum to initiate infection. Enhancing soil nitrogen content through cultural practices is also helpful to reduce severity of the disease. Chemical control can be achieved by preventive treatments with mancozeb, propiconazole, tebuconazole or azoxystrobin if sprayed at 10-day intervals (Rochecouste, 1984; Blum & Gabardo, 1993; Koike *et al.*, 2001). Planting of resistant cultivars is reported to be efficient, economical, environmentally friendly and simple approach for managing garlic rust. Rust have been controlled as part of an integrated crop management system using resistant cultivars and chemical control to complement the action of one with another although the disease is still severe in many parts of the world. This brief piece of work was, therefore, carried out with the objective to review the effect rust disease on garlic production and its management options.

Life cycle and Taxonomic Position of *Puccinia allii*

Puccinia allii is an autoecious rust, producing pycnia, aecia, uredinia, telia, and basidia on a single host (Anikster *et al.*, 2004). The genus *Puccinia* is categorized under the Kingdom-Fungi, Division/Phylum-Basidiomycota, Class- pucciniomycetes, Order-Uredinales and Family-Pucciniaceae. This genus is the largest in the order Uredinales (Anikster *et al.*, 2004). The taxonomic delineation of *Puccinia allii* has been in flux over the last century due to variation in the host range (Anikster *et al.*, 2004; Metcalf and Napier, 2002; Lupien *et al.*, 2004). Koike *et al.* (2001) and Anikster *et al.* (2004) determined that rust on garlic and chives in the USA was a different species to that on wild leek, leek and garlic in Europe, as leek was not susceptible to infection by the American rust following experimental inoculations. In addition, they found the two clades differed in the number of nuclei in their basidiospores, and by differences in the size of their teliospores (Anikster *et al.* 2004). The rust from the USA that has a majority of two-celled teliospores, and infects garlic and chives is hereafter referred to as *P. allii* (Koike *et al.*, 2001).

The current classification of *P. allii* is considered a species complex rather than a single species and includes *P. porri*, *P. mixta*, *Uromyces ambiguus*, and *P. blasdalei* and *U. duris*. At present two members of this species complex have been clearly defined and are represented by the North American collections from garlic and Middle Eastern collections from leek (Szabo *et al.*, 2008). Resolution of *puccinia allii* is complex will require further sampling of rusts on other species of *Allium* that can be linked to earlier described species (Alistair *et al.*, 2016).

Generally

Anikster *et al.* (2004) reported great differences in morphology between North American and European/Middle Eastern isolates, suggesting that this rust consists of at least two species.

Geographical Distribution and Economic Importance of Garlic Rust

Puccinia allii is a common cosmopolitan pathogen, present in onions, garlic and leeks, and reported in Europe, Asia, Africa, North and South America, Australia and New Zealand (Koike *et al.*, 2001; Anisker *et al.*, 2004; Fuyuru *et al.*, 2009; Alistair *et al.*, 2016). Rust on garlic, caused by the fungus *Puccinia allii*, has a worldwide distribution (Koike *et al.*, 2001); including Africa, Asia, Australasia, Europe, the Middle East, North and South America (Anikster *et al.*, 2004).

It is an important disease affecting leek crops as well as garlic, and occurs in the United Kingdom (Smith *et al.*, 2000) Netherlands (De Jong *et al.*, 1995) and USA (Koike *et al.*, 2001). In 1998, a devastating outbreak of rust disease affected garlic throughout California. The disease was widespread, and in many fields 100% of the plants were infected. It also occurred in 1999 and 2000, indicating that rust may have become an annual problem in some parts of the State (Koike *et al.*, 2001). It is also an important disease reported in different parts of Ethiopia (Mengesha *et al.*, 2016; Worku, 2017; Yeshwas *et al.*, 2018).

Worldwide, garlic rust has caused significant losses to garlic, leek, and onion production. The outbreak resulted in yield losses of 51% and an economic loss of 27% to the industry (Anikster *et al.*, 2004). In the late 1990s, outbreaks in the United States of America reduced crop yields by up to 75% in some fields (Janet and Tammy, 2008). Yield losses as high as 83% have also been reported by Ahmad and Iqbal (2002) on garlic due to the same disease in Nepal. Bulbs infected with garlic rust remain small and are of low storage quality (Sartaj and Ahmad, 2005). According to the authors this disease also causes considerable loss on different Allium crops. It is also an important disease in Ethiopia too as it resulted in yield losses of 48% (Worku and Dejene, 2012) and almost 50% (Mengesha *et al.*, 2016). Garlic rust is the most important disease of the crop in the highlands of Bale and causing a total bulb yield loss as high as 58.75% (Worku, 2017).

Means of Dispersal and Epidemiology of the Disease

The initiation of rust epidemics depends critically on rainfall that effectively deposits spores travelling long distances (Nagarajan and Singh, 1990). The first sign of infection in a spring-sown crop usually appears between June and August in the temperate regions. The fungus probably survives as either uredospores or teliospores, with uredospores apparently being the most important source of inoculum. Uredospores are wind borne and can be spread long distances. The cyclical production of uredospores during the growing season serves as a continuous inoculum source throughout the life cycle of the crop (Hill, 1995). Teliospores, a second type of spores formed by garlic rust fungus, later develop on the same leaves, resulting in black pustules (Timila *et al.*, 2005).

Symptoms

Puccinia allii infects garlic at bulb formation stage (Koike *et al.*, 2001). Garlic rust is readily identified by the earliest symptom of small, circular to elongate white flecks that occur on both sides of leaves, as the disease progresses, these small spots expand, and the leaf tissue covering the lesions ruptures and masses of orange, powdery spores (uredospores) become visible as pustules (Schwartz and Mohan, 2008; Worku, 2017). The pustules become full of uredospores that spread the disease to plants by rain splash and wind currents. Severely infected leaves are almost entirely covered with pustules, resulting in extensive yellowing, wilting and premature drying of leaves. Teliospores, a second type of spores formed by garlic rust fungus, later develop on the same leaves, resulting in black pustules (Koike *et al.*, 2001). The earliest severe rust on garlic and other Alliums can cause extensive loss of foliage and subsequent reduction in bulb size and quality. On infected onion and chives, symptoms consist of small (less than 3.2 mm in diameter), white to tan spots. The orange pustules often form concentric groups on the spot periphery. Disease severity on onion and chives is significantly less severe than on garlic. Generally, this garlic rust can cover entire leaves, causing them to die (Tahir *et al.*, 2006).

Host Plant Ranges and Pathogen Survival

Puccinia allii causes rust diseases in onion (*Allium cepa* L.), scallion (*A. chinense* L.) spring onion or Japanese bunching onion (*A. fistulosum* L., porrum group: naganegi), garlic (*A. sativum* L. nin-niku), and Chinese chive (*A. tuberosum* L.) in Japan (Foriaya *et al.*, 2009).

In Australia Metcalf (2002) showed in pathogenicity tests that an isolate of *Puccinia allii* from Tasmania could infect chives, garlic, bulb onions, spring onions and shallot. Severe infections produced by this species complex can kill leaves of cultivated species of *Allium*. Garlic, chives, and leeks are the most susceptible crops.

Koike et al. (2001) found no evidence for the disease being maintained or spread by volunteer plants or through an alternate host and concluded that the disease was carried from one season's crop to the next by a living bridge of infected plants, since there are crops in the field all year round; uredospore could therefore be blown from either local or more distant foci of infection. They also persist in the soil on plant debris for several years. Other kinds of spores in the complex life cycle of rusts are produced in other *Alliums*. It has been reported that the aforementioned pathogens also infect Welsh onions (*Allium fistulosum* L.), chives (*Allium schoenoprasum* L.) and other alliums (Szabo et al. 2008).

Management Options

Diseases caused by *Puccinia allii* are controlled primarily through the use of chemical sprays with fungicides, and disease free or treated seeds, adequate nitrogen fertilizer were generally reduces the rate of infection by *Puccinia allii*. Crop rotation, removal and burning of crop debris (if infected), and eradication of weed hosts help in reducing the inoculum potential for subsequent plantings of susceptible crops (Agrios, 2005). For management of garlic rust, the following options have been worked out.

Cultural control

Crop health is affected by cultural practices such as planting time, plant density, cropping sequence, cropping pattern, fertilization, seed-bed preparation, weeding, etc. Such practices can also influence the level of biotic stresses and their effect on the growth and yield of a crop. Protection often provided by modification the environment to favor crop growth, but discourage the conditions that favor disease development. Cultural methods, which can be manipulated to control disease, include use of healthy planting materials, fertilizer application, crop rotation, optimum planting date, supplying adequate moisture and optimum planting density (Agrios, 2005).

Use of healthy planting materials

Puccinia allii can survive on the bulb of garlic. Hence, use of healthy seed can reduce the severity of the disease on the developing crop (Greisbach and Ocamba, 2009; Janet and Tammy, 2008). Garlic rust pathogen is a fungal microorganism that survives either on crop debris by producing resting spores or alternates with different, perennial hosts, like wild *Allium* species. Hence, rotating out of *Allium* crops for two to three years and destroying all volunteers and *Allium* weeds can be effective in reducing the amount of inoculum in the crop residue in the soil for the next growing season (Schwartz and Mohan, 2008; Greisbach and Ocamba, 2009).

Adequate supply of moisture

The garlic rust occurs most frequently under conditions of high humidity and low rainfall and its incidence is highest in stressed plants (Janet and Tammy, 2008). It is also reported that the disease can cause severe yield losses when excessive rain, fog, or irrigation are present (Greisbach and Ocamb, 2009). According to these authors supply of optimum moisture to the crop is important to reduce garlic rust problem. Garlic rust is highly dependent on moisture, changes in precipitation or dew will directly influence the severity and distribution of the disease.

Optimum plant density

Mohibullah (1991) reported that rust severity could be decreased by 10% by decreasing plant population from 2 to 0.5 million per hectare and consequently garlic yield could be increased by 28%. The maximum value of yield was obtained from plot planted with 10 cm intra row spacing while, the minimize value of yield was achieved by plot planted with 20 intra row spacing at research field of haramaya university, Ethiopia (Mengesha, 2015). A number of studies in various parts of the world have shown that garlic production can be improved through proper spacing that plant spacing significantly increase number of cloves per bulb, bulb size, bulb weight and yield. It has been reported by (Mohammad et al, 2014). Naruka and Dhaka (2001) and Alam, et al. (2010) indicated that garlic bulb yields increased significantly with increasing intra-row spacings. It is assumed that densely populated plants are ideal habitats for fungal development and transmission to the nearby plant since moisture is retained within the leaves, and it prevents direct sunshine (Jorind, 2012). Therefore, increasing spacing between garlic plants will

allow better airflow and keep humidity from building up. Increasing plant spacing can significantly decrease disease incidence and severity level of garlic rust. This might be due to proper aeration resulting in decreased humidity level suitable for fungus growth and decreasing plant population also limits the transmission of rust pathogen to the next plant (Ahmed *et al.*; 2017).

Fertilizer application

Nitrogen (N) is essential for plant growth and development. In addition nitrogen availability can influence the resistance of plants to plant pathogens (Marschner, 1995, Tadesse and Mashila, 2018). Too much nitrogen and too little potash in the soil would promote rate of infection. Lush garlic growth caused by excessive nitrogen will make the crop susceptible to rust infection (Greisbach and Ocamba, 2009). Generally, adequate nitrogen fertilizer reduces the rate of infection (Agrios, 2005).

Host Resistant

Breeding for genetic resistance is one of the best methods to handle fungal diseases, as it is the most economical and environment-friendly control method. However, Koike *et al.* (2001) tested 34 garlic varieties against garlic rust and noted that acceptable levels of resistance were not observed in any cultivar. Again Yeshewas *et al.* (2018) screened 16 garlic germplasm for yield and diseases tolerance in east Gojam and reported none of the germplasm showed high resistance to rust. Additionally use of resistant varieties is cheaper but such varieties are not available (Worku, 2017). Because garlic is a sterile plant that is propagated asexually by way of vegetative cloves, bulbs, and bulbils, not from seed, which markedly reduces chances for genetic improvement (Aparicio *et al.*, 2011; Kamenetsky and Rabinowitch, 2017). However, recent efforts have shown that fertility restoration is possible (Pooler and Simon, 1994; Kamenetsky *et al.*, 2005) allowing genetic studies and classical breeding.

Biological control

Biological control is currently receiving much attention in disease management practices. The over uses of chemicals has created an unprecedented environmental crisis. It is defined as brings about reduction in activities of pathogens by other organisms and typically involves an active human role. The majority of work done on plant disease bio-control was related to soil-borne diseases using either bacteria or fungal antagonists. However, there is no more published source available regarding biological control against garlic rust on garlic (Tahir *et al.*, 2006). In Ethiopia commercial biological control is not available for garlic rust (Worku, 2017).

Chemical control

There are different recommended fungicides that are registered for the control of garlic rust on different *Allium* crops in different countries. Chemical control can be achieved by preventive treatments with mancozeb, propiconazole, tebuconazole or azoxystrobin, if sprayed at 10-day intervals (Koike *et al.*, 2001). Four chemical fungicides were evaluated for their efficacy against garlic rust at Hararge highlands. There was significant garlic rust severity difference between fungicide-treated and the unsprayed checks. garlic plots treated with propiconazol showed the lowest disease severity (3.72%) whereas plots treated with bayleton showed greater disease severity (73.47%) which is nearest to the control (83.45%) at the 128 days (Mengesha *et al.*, 2016). With the Section 18 registration for tebuconazole, commercial garlic was sufficiently protected during the 1999 and 2000 seasons (Koike *et al.*, 2001). All fungicide treated plots gave higher total yield than the unsprayed check plots (Worku, 2012). This author also reported early garlic rust detection and application of fungicide while the disease was less severe is very essential if effective control measure is intended. Using systemic fungicides contains tebuconazole as active ingredient recommended in different countries for the management of garlic rust and other disease of garlic in different countries (Maria *et al.*, 1998, Koike *et al.*, 2003, Zewde *et al.*, 2007). Garlic rust (*Puccinia allii*) severity level on garlic plots sprayed with the three different Nativo SC 300(Trifloxystrobin 100g/l +Tebuconazole 200g/l) spray frequencies (five times in every 7-days, three times in every 14-days and two times in every 21-days) was lower as compared to the unsprayed plots (Worku, 2017). In addition Marketable yield harvested from sprayed plots was consistently greater than the yield harvested from unsprayed plots (Tilahun *et al.*, 2018).

Integrated Disease Management

To reduce yield loss of garlic due to rust integrating all components-cultural practices, host resistant, chemical control, biological control, and optimum crop density into a disease management scheme that are environmentally compatible, economically feasible, and socially acceptable to reduce damage caused by disease to tolerable levels. Integrated disease management (IDM) is a combination of methods such as cultural, biological and host resistance that are environmentally compatible, economically feasible, and socially acceptable to reduce damage caused by disease to tolerable levels. The effect of *Puccinia allii* on garlic can be minimized by integrated use of different control measures.

References

- Agrios, G.N. (2005). Plant Pathology, 5th Ed. Academic Press, San Diego. 922p.
- Alam M., Rahim M., Bhuyan MAH, Simonand P., Malek M (2010). Effect of spacing on growth and yield of two lines of garlic under dry land condition. *Journal of Agroforestry and Environmental science*, 4: 151-154.
- Anikster, Y., Szabo, L.J., Eilam, T., Manisterski, J., Koike, S.T., and Bushnell, W.R (2004). Morphology, life cycle, biology and DNA sequence analysis of rust fungi on garlic and chives from California. *Phytopathology*, 94: 569 -577.
- Blum L.E.B., Gabardo H. (1993) Chemical control of the garlic rust in Curitiba/SC, Brazil. *Fitopatologia Brasileira*, 18: 230–232.
- Chia-Wen T., C.Haw-Wen, S.Le-Yen, L.Chong-Kuei (2012). Garlic: Health benefits and actions. *Biomedicine* 2: 17-29.
- CSA (Central Statistical Agency). (2017). The federal democratic republic of Ethiopia central statistical agency agricultural sample survey (Meher season, private peasant holding), Statistical Bulletin, Number, 584. Addis Ababa, Ethiopia.
- FAO (Food and Agriculture Organization). (2012). Area and production of crops by countries.
- Foriya, H., Takanashi, H., Fuji, S., Nagai, Y., and Naito, H. (2009). Modeling infection of spring onion by *Puccinia allii* in response to temperature and leaf wetness. *Phytopathology*, 99:951-956.
- Figliuolo, G., Candido, V., Logozzo, G., Miccoli, V., and Spagnoletti-Zeuli, P.L., (2001). Genetic evaluation of cultivated garlic germplasm (*Allium sativum* L. and *Allium ampeloprasum* L). *Euphytica*, 121:325-334.
- De Jong, P.D., Daamen, R.A., and Rabbinge, R., (1995). The reduction of chemical control of leek rust, a simulation study. *European Journal of Plant Pathology*, 101: 687 – 693.
- Getachew, T. and Asfaw, Z. (2000). Achievements in shallot and garlic research, EARO, Addis Ababa, Ethiopia.
- Greisbach, J. and Ocamb, C. M. (2009). Garlic (*Allium sativum*) rust. Online guide to plant disease control. Arizona State University. USA.
- Hill, J.P. (1995). Rust. Pp. 24-25. In: Compendium of onion and garlic diseases. Schwartz HF, Mohan SK, (eds). St. Paul, Minnesota: APS Press.
- Janet, B., and Tammy, H., (2008). Garlic: Organic production The epidemiology of plant diseases. Kluwer Academic Publishers, UK, 460 pp.
- Jennings, M., Brian, V., Ford-Lloyd and Gillian, M., 1990. Rust infections of some *Allium* species: An assessment of germplasm for utilizable rust resistance. *Euphytica*, 49: 99-109.
- Jong, P., (1995). Growth of leek rust epidemics in time in three cultivars during the early stage of the epidemic. *European Journal of Plant Pathology*, 101: 139-148.
- Kamenetsky R., London Shafir I., Khassanov F., Kik C., Van Heusden A.W., Vrieling-Van Ginkel M., Burger-Meijer K., Auger J., Arnault I., Rabinowitch H.D. (2005). Diversity in fertility potential and organo-sulphur compounds among garlics from Central Asia. *Biodiversity and Conservation* 14: 281–295.
- Koike, S.T., Smith, R.F., Davis, R.M., Nunez, J.J. and Voss, R.E. (2001). Characterization and control of garlic rust in California. *Plant Disease* 85: 585-591.
- Lea, P.J. and Azevedo, R.A. (2006). Nitrogen use efficiency. uptake of nitrogen from the soil. *Annals of Applied Biology* 149: 243 - 247.
- Lea, P.J., Sodek, L., Parry, M.J., Shewry, P.R. and Halford, N.G. (2007). Asparagine in plants. *Annals of Applied Biology* 150: 1 - 26.
- Libner, N. (1989). Vegetable production. Van Nostrand Rein hold, New York, USA, pp. 657.

- Lupien, S.L., Hellier, B.C. and Dugan, F.M. 2004. First report of onion rust caused by *Puccinia allii* on *Allium pskemense* and *A. altaicum*. *Plant Disease* 88: 78-83.
- Mengesha, W., Tesfaye, A. and Mashilla, D. (2015). Effect of spacing in incidence and severity of garlic rust (*Puccinia allii*) and bulb yield and related traits of garlic at eastern Ethiopia. *Plant Pathology and Microbiology* 6: 314-338.
- Mengesha, W., Azene, T. and Mashilla, D. (2016). Evaluation of fungicides on the control of garlic rust (*Puccinia allii*) in Eastern Ethiopia. *International Journal of Emerging Technology and Advanced Engineering* 6: 27-33.
- Metcalf, D.A., and Napier, T (2002). Host range of Tasmanian strains of onion rust. In: *Proceedings of Onions Conference, National Vegetable Industry Centre, Yanco Agricultural Institute, Australia*. 72pp.
- Mohibullah, K (1991). Studies on major diseases of bulb vegetables (onion and garlic) in (NWFP). Final Technical Report Agricultural Research Institute Tarnab, Peshawar.
- Nagarajan, S. and Singh, D.V (1990). Long-distance dispersion of rust pathogens. *Annual Review of Phytopathology* 28: 139 - 154.
- Negash, T., Shifa H., Regassa, T. (2018). Management of Garlic Rust (*Puccinia allii*) Through Fungicide at Bale Highlands, South Eastern Ethiopia. *Food Science and Quality Management* 81: 80-85.
- Pooler M.R., Simon P.W. (1994) True seed production in garlic. *Sexual Plant Reproduction*, 7, 282–286.
- Rochecouste J.F.G. (1984). Chemical control of garlic rust. *Australasian Plant Pathology* 13: 47–48.
- Sartaj, S. and Ahmad, S (2005). Identification of rust pathogen species isolated from different garlic cultivars planted in district Swabi, NWFP. *Sarhad Journal of Agriculture* 21: 205-213.
- Smith, B. M. and Crowther, T. C (1992). Field assessment of the reaction of leek cultivars to leek rust. *Annals of Applied Biology*, 120: 104 - 105
- Schwartz, H.F., Mohan, K., Havey, M. and Crowe, F. (2008). The genus *Allium*. pp. 1-6. In: *Compendium of onion and garlic diseases*. H.F. Schwartz and S.K. Mohan. American Phytopathological Society (APS). St. Paul, Minnesota, USA.
- Szabo, L.J., Koike, S.T. and Hill, J.P (2008). Rust. In: Schwartz, H.F, and Mohan, S.K. *Compendium of Onion and Garlic Diseases and Pests*. 2nd edition. 44pp.
- Tahir M, Ahmad M, Shah M, Alam S, Khattak M (2006). Field efficacy of different spray fungicides on the severity of garlic rust (*Puccinia porrii* Wint). *Sarhad Journal of Agriculture* 22: 237 - 241.
- Tesfaye, T. and Habtu, A. (2003). A review of vegetable diseases in Ethiopia. pp. 495-518. In: *A Review of Crop Protection Research in Ethiopia*. Tsedeke Abate (Ed.). Department of Crop Protection, Institute of Agricultural Research, Addis Ababa.
- Tadesse, M., Dejene, M. (2018). Effect of Nitrogen and Fungicidal Spray Rates on Incidence and Severity of Garlic Rust (*Puccinia allii*) at Haramaya, Ethiopia. *Advances in Life Science and Technology* 63: 24-29.
- Tindall, H.D. (1983). *Vegetables in the tropics*. The Macmillan press Ltd, Hong Kong. pp.533.
- Timila R., Joshi, S., Manandhar, G., Sharma, S. (2005). Occurrence of garlic rust in Mid Hill of Nepal. *Nepal Agricultural Research Journal*, 6: 110-111.
- Toome, M., Heinso, K., Holm B., Luik A. (2011). The influence of canopy density on willow leaf rust (*Melampsora epitea*) severity in willow short rotation coppice. *Biomass and bioenergy*, 34: 1201-1206.
- White, J., Ryley, M. George, G., Kong, A. (2014). Optimal environmental conditions for infection and development of *Puccinia purpurea* on sorghum. *Australasian Journal of Plant Pathology* 43: 447-457.
- Worku, Y. and Dejene, M. (2012). Effects of garlic rust (*Puccinia allii*) on yield and yield components of garlic in Bale highlands, south eastern Ethiopia. *Journal of Plant Pathology and Microbiology* 3:218-2224.
- Worku, Y. (2017). Determination of optimum native SC 300 (Trifloxystrobin 100g/l + Tebuconazole 200g/l) spray frequency for control of rust (*Puccinia allii* Rudolphi) on garlic in Bale highlands, south eastern Ethiopia. *American Journal of Agriculture and Forestry* 5: 16-19.
- Yeshiwas.Y, Belete N., Tegibew, W., Yohaness, G., Abayneh, M., and Kassahun, Y (2018). Collection and characterization of garlic (*Allium sativum* L.) germplasm for growth and bulb yield at Debre Markos, Ethiopia. *Journal of Horticulture and Forestry* 10:17-26.
- Zewide, T., Fininsa, C. and Sakhuja, P. (2007). Management of white rot (*Sclerotium cepivorum*) of garlic using fungicides in Ethiopia. *Crop Protection* 26: 856–866.